



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

come from the brown algae, and even the Characeae showing more resemblance to the brown algae than to other green algae. The origin of antheridia and archegonia from a plurilocular sporangium is developed along the lines already presented by DAVIS and HOLFERTY. To some of us, it would seem better to derive antheridia and archegonia from plurilocular sporangia of some hypothetical green alga than to refer them directly to the plurilocular sporangia of brown algae. The spore mother cells of archegoniates are compared with the unilocular sporangia of the brown algae, and the sporophyte of archegoniates with the thallus of brown algae. SCHENCK does not believe the sporophyte of pteridophytes can be derived from that of bryophytes. Even the complicated antheridium of the Characeae is referred to the plurilocular sporangium of the brown algae.—CHARLES J. CHAMBERLAIN.

Translocation in green tissues.—RYWOSCH points out²² that translocation must depend upon the concentration gradient from the peripheral cells to the vascular bundle. This gradient is due in part to the excess of food made in the cells best illuminated, and also to the fact that transpiration cooperates doubly, by reducing the amount of water and by determining the movement of water. Thus those cells next the bundle are first to receive the water supply and those nearer the periphery are driest. He shows that the emptying of leaf tissues is not simultaneous, peripheral ones being emptied first, and that the whole process is greatly retarded when transpiration is checked. [Yet it must not be forgotten that there are plants in which transpiration cannot be invoked as an aid to translocation, since it is practically non-existent for weeks or months at a stretch.] The concentration is also kept low in the inner cells by the making of starch in them. RYWOSCH also adds a note on the function of the starch sheath, holding that its character as a reserve is very doubtful.—C. R. B.

Hygroscopic living leaves.—HANNIG²³ reports what he says is the first recorded instance of the movement of living leaves produced by variations in the water-content of the cell walls. The leaves of various hardy species of *Rhododendron* rise and fall, roll and unroll, according as they are subject to freezing and thawing weather respectively, though the same movements may be produced by other conditions which reduce or increase the water-content of the cell walls. Turgor is not concerned, HANNIG says, because dead or live, narcotized or unnarcotized, leaves exhibit movements equal in extent and kind. HANNIG's argument is not convincing, and it seems unlikely that this conclusion is sound. In fact, too little is known of the physics of water and cell contents under the conditions described to make it possible to state accurately the precise relations involved.—C. R. B.

²² RYWOSCH, S., Sur Stoffwanderung im Chlorophyllgewebe. Bot. Zeit. **66**: 121-130. figs. 2. 1908.

²³ HANNIG, E., Ueber hygroscopische Bewegungen lebender Blätter bei Eintritt von Frost und Tauwetter. Ber. Deutsch. Bot. Gessells. **26a**: 151-166. 1908.